



Utilization Management Policy

Scoliosis: Conservative Interventions

Table of Contents		Related Policies	Policy Number	95
Policy Statement.....	1	Experimental and Investigational Services and Devices	Original Effective Date:	1/1997
Purpose.....	1	General Guidelines for use of Plain Film Radiography	Current Approval Date:	4/27/2023
Scope.....	1		Next Review:	4/2024
Background.....	2		Category:	Determination
Literature Review.....	4			
References.....	13			
History.....	15			
Plain Language				
Summary.....	16			

Policy Statement

Optum* by OptumHealth Care Solutions, LLC considers brace therapy that utilizes a rigid orthosis (e.g., Boston Brace) to be proven and medically necessary for the sustained reduction and/or stabilization of curve magnitude, when patient selection criteria have been satisfied.

Optum considers manual therapy (including manipulation), exercise (including specific exercise approaches), soft bracing (e.g., SpineCor), whole body vibration, and non-operative traction therapies to be unproven and not medically necessary for the sustained reduction and/or stabilization of curve magnitude due to insufficient scientific evidence of efficacy for the treatment of idiopathic scoliosis.

The use of manual therapy, exercise and soft braces for the treatment of idiopathic scoliosis deformity is supported by some positive published information regarding safety and/or efficacy. However, a beneficial impact on health outcomes (e.g., durable curve reduction) has not been proven because the data are sparse, and the evidence is of very low-quality.

The research evidence regarding the use of whole-body vibration and non-operative traction therapy in the treatment of idiopathic scoliosis is so limited that an appraisal of safety and efficacy cannot be made.

Purpose

This policy has been developed as the clinical criterion that describes the position of Optum regarding the efficacy, effectiveness, risks, and burdens associated with the use of conservative interventions (manual therapy, exercise, bracing, whole body vibration and non-operative traction) for the treatment of idiopathic scoliosis.

Scope

This policy applies to all in and out of network programs, involving all provider types, where utilization review (UR) determinations are rendered. This policy serves as a resource for peer-to-peer interactions in describing the position of Optum on the clinical appropriateness and/or medical necessity of conservative interventions for the treatment of idiopathic scoliosis

*Optum™ Physical Health (“Optum”) includes OptumHealth Care Solutions, LLC; ACN Group IPA of New York, Inc.; ACN Group IPA of California, Inc. d/b/a OptumHealth Physical Health of California; Managed Physical Network, Inc.; and OrthoNet Holdings, Inc. which includes OrthoNet New York IPA, Inc., OrthoNet West, Inc., OrthoNet, LLC, OrthoNet of the South, Inc.

The application of this policy is limited to the conservative (non-operative) treatment of idiopathic scoliosis for the purpose of any of the following goals: arresting curve progression, slowing curve progression, or reducing the magnitude of curvature. Conservative interventions included in this policy encompass manual therapy, exercise, bracing, whole body vibration, and non-operative traction.

Key Questions

1. Is there sufficient research evidence of the efficacy and safety of manual therapies e.g., manipulation, mobilization, soft-tissue techniques, etc. in the treatment of idiopathic scoliosis for the sustained reduction and/or stabilization of curve magnitude to conclude this/these interventions are an appropriate therapeutic alternative for patients diagnosed as having idiopathic scoliosis?
2. Is there sufficient research evidence of the efficacy and safety of exercise therapy in the treatment of idiopathic scoliosis for the sustained reduction and/or stabilization of curve magnitude to conclude this/these interventions are an appropriate therapeutic alternative for patients diagnosed as having idiopathic scoliosis?
3. Is there sufficient research evidence of the efficacy and safety of rigid and/or soft brace therapy in the treatment of idiopathic scoliosis for the sustained reduction and/or stabilization of curve magnitude to conclude this/these interventions are an appropriate therapeutic alternative for patients diagnosed as having idiopathic scoliosis?
4. Is there sufficient research evidence of the efficacy and safety of whole-body vibration in the treatment of idiopathic scoliosis for the sustained reduction and/or stabilization of curve magnitude to conclude this intervention is an appropriate therapeutic alternative for patients diagnosed as having idiopathic scoliosis?
5. Is there sufficient research evidence of the efficacy and safety of non-operative traction therapy in the treatment of idiopathic scoliosis for the sustained reduction and/or stabilization of curve magnitude to conclude this intervention is an appropriate therapeutic alternative for patients diagnosed as having idiopathic scoliosis?

Background

Overview:

Idiopathic scoliosis is defined radiographically as a lateral curvature of the spine greater than or equal to 10° Cobb with rotation, of unknown etiology.¹ Idiopathic scoliosis is most commonly identified (~90% of cases) during adolescence (ages 10 – 18 years).² Idiopathic scoliosis progresses most often in adolescents who are growing and have curves which are above 20 degrees. This is the time conservative interventions including bracing are commonly considered.³ The efficacy of conservative treatment in adolescent idiopathic scoliosis (AIS) is, however, controversial due to variations in inclusion and assessment criteria⁴, as well as sparse and low-quality evidence.⁵

Indications for Conservative Intervention:

The primary aim of scoliosis management is to stop curvature progression.⁶⁻⁸ Guidance for intervention is broadly based on the risk for significant curvature progression in a given period of time.^{3,6,7} Rowe described decision-making about treatment strategies by assessing a combination of Risser Sign (skeletal immaturity) and Cobb angle (curve magnitude) measurements. [Table 1]

*Optum™ Physical Health (“Optum”) includes OptumHealth Care Solutions, LLC; ACN Group IPA of New York, Inc.; ACN Group IPA of California, Inc. d/b/a OptumHealth Physical Health of California; Managed Physical Network, Inc.; and OrthoNet Holdings, Inc. which includes OrthoNet New York IPA, Inc., OrthoNet West, Inc., OrthoNet, LLC, OrthoNet of the South, Inc.

Table 1: Indications for Treatment

Risser Sign	Cobb Angle (degrees)	Action
0 – 1	0 – 20	Observation
0 – 1	20 – 40	Bracing
2 – 3	0 – 30	Observation
2 – 3	30 – 40	Bracing
0 – 3	40 – 50	Bracing
0 – 4	> 50	Surgery

Adapted from: Rowe DE. The Scoliosis Research Society brace manual: introduction. *Scoliosis Research Society* 2003

The Society on Scoliosis Orthopaedic and Rehabilitation Treatment (SOSORT) has published guidelines on the indications for the conservative management of scoliosis.^{8,9} The guidelines are intended to apply to all idiopathic scoliosis patients regardless of age. The main clinical questions that they cover are:

- Which assessment of the patient should be performed?
- Which conservative treatment should be provided and how?
- How and when should bracing be applied?
- How and when should exercises be used?

The 2016 version of SOSORT guidelines include a strength of treatments scheme (STS), which stratifies recommendations by curve severity, age, Risser stage, and the presence of pain and/or trunk decompensation.⁸

There were several limitations associated with the 2016 SOSORT guidelines, which mitigate their ability to inform policy. The guidelines employed a parsimonious strength of evidence scheme, based on study design and numbers of studies. The strength of recommendation reflected the relative importance of the recommendation and how broadly the recommendation applied, as opposed to the extent to which a recommendation is likely to be affected by new evidence. The literature review, while comprehensive, was reported in a narrative format. The quality of the evidence (ie, confidence in the estimates of effect) was not evaluated. Further, the review did not evaluate important considerations of the body of evidence such as precision, consistency, and risks as part of the evidence review.

Broadly, scoliosis-specific exercises are recommended as the first step to treat idiopathic scoliosis to prevent/limit progression of the deformity and bracing. The SOSORT recommendation for exercise to impact curve progression was described as, "...less important, it can be applied on a voluntary basis."⁸ This recommendation is based upon the findings of multiple randomized controlled trials (RCTs) or systematic reviews of RCTs.

The nine references supporting this recommendation are listed in Table 11 of the SOSORT guideline.⁸ An assessment of these sources showed there was single RCT that investigated effectiveness of exercise on curve progression.¹⁰ This study was included as part of a systematic review¹¹, which has been incorporated into the policy's literature review. Another reference was a systematic review that is also as part of the policy literature review.¹² A third study reported as a reference in the SOSORT guideline was a preliminary cohort design (N=42) that investigated effectiveness of exercise on curve progression over a 3-month period.¹³ None of the other studies cited by the SOSORT guidelines evaluated the effect of exercise on curve progression for individuals diagnosed with idiopathic scoliosis. [Table 2]

*Optum™ Physical Health ("Optum") includes OptumHealth Care Solutions, LLC; ACN Group IPA of New York, Inc.; ACN Group IPA of California, Inc. d/b/a OptumHealth Physical Health of California; Managed Physical Network, Inc.; and OrthoNet Holdings, Inc. which includes OrthoNet New York IPA, Inc., OrthoNet West, Inc., OrthoNet, LLC, OrthoNet of the South, Inc.

Table 2: Sources cited to support SOSORT (2016) recommendation for scoliosis-specific exercises as the first step to treat idiopathic scoliosis to prevent/limit progression of the deformity and bracing

Author (Date)	Study type	Investigated effectiveness of exercise on curve progression?	Notes
Monticone ¹⁰ (2014)	RCT	Yes	Included in appraised systematic review(s)
Wong ¹⁴ (2005)	Cohort	No	The study compared two methods of spinal cast/modelling.
Périé ¹⁵ (2004)	Laboratory study	No	The study analyzed patient-specific bracing biomechanics in the treatment of scoliosis
Weiss ¹⁶ (2006)	Consensus report	No	Report describes a baseline for developing a consensus for language and goals for proposed multicenter clinical studies regarding scoliosis.
Lenssinck ¹² (2005)	Systematic review	Yes	Included in literature review
Stone ¹³ (1979)	Cohort	Yes	Preliminary study
Diab ¹⁷ (2012)	RCT	No	The study investigated the effectiveness of forward head correction on postural parameters and functional level in AIS patients.
Toledo ¹⁸ (2011)	RCT	Yes	Not idiopathic scoliosis. The study participants were diagnosed with nonstructural (functional) scoliosis.
McMaster ¹⁹ (1983)	Case series	No	This study reported on the incidence of idiopathic scoliosis.

The 2016 SOSORT guidelines recommended bracing to treat adolescent idiopathic scoliosis (important but does not have to be applied to all patients with a specific need; based on multiple RCTs or systematic reviews of RCTs).⁸

Literature Review

Search Strategy:

An updated literature search was conducted on January 24, 2022. A comprehensive literature search was conducted using guidance provided by the Cochrane Handbook for Systematic Reviews of Interventions.²⁰ Biomedical databases and consumer-oriented search engines were used to identify and retrieve relevant evidence. Hand-searches of bibliographies and non-indexed documents were included in the search strategy. Additionally, professional specialty society websites e.g., SOSORT (<http://sosort2014.com/>), Scoliosis Research Society (<http://www.srs.org/>) were searched for research evidence. Research in-progress and protocols were identified by searching www.clinicaltrials.gov and published protocols.

Inclusion/Exclusion Criteria

Studies were included if they were designed to report on the effectiveness of one or more conservative interventions for the treatment of persons having been diagnosed with idiopathic scoliosis, and the intervention goal was to reduce, arrest or slow down the progression of curvature as measured by Cobb angle. Studies not meeting these criteria were excluded. Additionally, studies were excluded from independent appraisal if they were already evaluated in at least one included systematic review. Evidence

*Optum™ Physical Health (“Optum”) includes OptumHealth Care Solutions, LLC; ACN Group IPA of New York, Inc.; ACN Group IPA of California, Inc. d/b/a OptumHealth Physical Health of California; Managed Physical Network, Inc.; and OrthoNet Holdings, Inc. which includes OrthoNet New York IPA, Inc., OrthoNet West, Inc., OrthoNet, LLC, OrthoNet of the South, Inc.

syntheses that did not employ qualitative methods (eg, narrative reviews) were not included in the evidence appraisal but may have provided background information.

Evidence Extraction:

Seven recently published evidence syntheses that evaluated exercise approaches and one overview of systematic reviews were identified. There were five relevant systematic reviews and two narrative reviews of manual therapy. Four systematic reviews for bracing that were included in the evidence appraisal. Previous versions of systematic reviews on the same topic by the same authors were not extracted for appraisal. A descriptive study regarding a form of non-operative traction (axial spinal unloading) was located in an earlier literature search. The previous literature search identified a single laboratory study concerning the effects of vibration on scoliotic spines. There were no clinical studies (observational or experimental) identified concerning the treatment of idiopathic scoliosis deformity with whole body vibration.

Evidence Appraisal:

Systematic reviews of bracing and manual therapy were critically appraised for quality using the AMSTAR (A MeaSurement Tool to Assess Reviews) instrument, which was developed to evaluate systematic reviews of randomized trials.^{21,22} The AMSTAR tool is comprised of 11 items that question the methodological quality of systematic reviews. It has good face and content validity. Literature reviews were rated as being methodologically strong (i.e., high-quality), if the AMSTAR score was ≥ 6 . This interpretation is consistent with the approaches of other ‘reviews of reviews’.²³⁻²⁵ The AMSTAR 2 tool was used to assess the quality of reviews of exercise therapy. The AMSTAR 2 tool has been used to assess risk of bias and the overall quality of reviews that include randomized and/or non-randomized studies of healthcare interventions. The AMSTAR 2 has 16 items in total, which includes 10 of the original 11 AMSTAR domains. It is not designed to generate an overall ‘score’. The overall rating scheme is based on weaknesses in critical domains that may greatly weaken the confidence that can be placed in a systematic review.²⁶

Five systematic reviews reported on manual therapy including manipulation for the treatment of idiopathic scoliosis [Table 3]. Three of the reviews were appraised as high-quality.²⁷⁻²⁹ All the reviews arrived at similar conclusions; there is a lack of evidence, which does not permit conclusions on the efficacy of manual therapy for the treatment of adolescent and adult idiopathic scoliosis. Two narrative reviews, not suitable for quality appraisal, critically evaluated the body of evidence concerning the efficacy of manual therapy for idiopathic scoliosis.^{30,31} Both reviews concluded that the evidence is insufficient to make judgments about the usefulness of various manual therapies in the treatment of idiopathic scoliosis.

For exercise as an intervention for the treatment of idiopathic scoliosis deformity, six systematic reviews and one meta-analysis were identified with the most recent literature search. [Table 4]. Overall, the findings of these evidence syntheses do not support any form of exercise as proven effective for the treatment of scoliotic deformities.

Two reviews were rated as moderate quality. In a systematic review of RCTs and non-randomized studies of an intervention (NRSI), Fan, et al. (2020) evaluated the effect of scoliosis-specific exercise (SSE) on scoliotic deformity improvement.³² The review included 10 studies encompassing 494 participants. SSE was compared to traditional exercise, standard care (observation or bracing) or any other non-SSE. The results, based on moderate quality evidence, determined there was insufficient evidence to prove that SSE with or without other conservative treatments can reduce Cobb angle or the angle of trunk rotation (ATR). Thompson, et al (2019) conducted a systematic review and meta-analysis of 9 RCTs (N=480) to assess the effectiveness of SSE on adolescent idiopathic scoliosis (AIS) compared with other non-surgical

*Optum™ Physical Health (“Optum”) includes OptumHealth Care Solutions, LLC; ACN Group IPA of New York, Inc.; ACN Group IPA of California, Inc. d/b/a OptumHealth Physical Health of California; Managed Physical Network, Inc.; and OrthoNet Holdings, Inc. which includes OrthoNet New York IPA, Inc., OrthoNet West, Inc., OrthoNet, LLC, OrthoNet of the South, Inc.

interventions.³³ The reviewers found only very-low-quality evidence to support the use of SSE rather than standard care or other types of exercise for patients with AIS to reduce spinal curvature and trunk rotation, improve pain, function, satisfaction with management and overall quality of life, although it is unclear if the reported differences are clinically relevant.

Five evidence syntheses were rated as critically low- to low-quality due to methodological flaws, limitations within the included primary studies, sparse evidence, imprecise and inconsistent results, indirectness, and an increased risk of publication bias. Burger, et al. (2019) found that Schroth exercises had a statistically significant effect on reducing the Cobb angle in adolescents with idiopathic scoliosis; however, the clinical relevance was not reported.³⁴ This review's findings should be considered with caution for physiotherapy practice because of the limited number of identified articles and their methodologic limitations. Day, et al. (2019) concluded, "There is insufficient evidence to suggest that both Schroth and SEAS (scientific exercise approach to scoliosis) methods can effectively improve Cobb angles in patients with AIS compared to no intervention. There is limited evidence that the SEAS method is more effective at reducing Cobb angles compared to traditional exercises in treating AIS."³⁵ Farooqui, et al. (2018) suggested that therapeutic exercise regimes alone have a pivotal role in both decelerating the progression of the curve and reducing the already increased magnitude of the curve.³⁶ However, the meta-analysis has more than one critical flaw and should not be relied on to provide an accurate and comprehensive summary of the available studies. Laita, et al. (2018) described the positive effects of therapeutic exercise based on the Schroth method or stabilization exercises.³⁷ However, it was not possible to describe the ideal moment for the intervention or the number of weekly sessions and the duration of each session. Gámiz-Bermúdez, et al. (2021) performed a systematic review and meta-analysis of 8 RCTs (N=279) that analyzed the efficacy of corrective exercise compared to other active and inert interventions in the improvement of deformity in AIS.³⁸ Overall, the reviewers described moderate-quality evidence for a medium effect (SMD = -0.52, 95% CI -0.96 to -0.1), favoring corrective exercise-based therapy for spinal deformity reduction. A reappraisal of the primary studies found the results were based on very low- to low-quality evidence – due to study limitations (RoB), inconsistency and imprecision – indicating that any estimate of effects is very uncertain, and that further research is very likely to have an important impact on our confidence in the estimate of effects and is likely to change the estimate. Additionally, the clinical relevance of the results could not be ascertained due to the method employed to assess effect size. Li, et al. (2021) systematically reviewed and meta-analyzed 9 studies (7 RCTs and 2 NRSI) involving 325 participants, assessing the effectiveness of core-based exercise for correcting a spinal deformity in individuals irrespective of age with scoliosis.³⁹ Based on very low-quality evidence, the exercise group had significantly lower Cobb angles (MD = -2.08, 95% CI: -3.89 to -0.28, P = 0.02). The difference was not, however, clinically significant (change of $\geq 5^\circ$). There was no significant difference observed regarding the angle of trunk rotation between groups (MD = -0.69, 95% CI: -2.61 to 1.22, P = 0.48).

The conclusion, the current body of evidence is insufficient to establish exercise therapy as proven effective for the treatment of scoliotic deformity, has been supported by an overview of systematic reviews and meta-analyses.⁴⁰ Zhou, et al. (2022) found exercise therapy to have potential benefits to treat patients diagnosed with AIS. However, the findings were not conclusive; given that some reviews relied on data from the trials with potential risk of bias and significant heterogeneity. More high-quality research is still needed to verify these findings.

Four of the appraised systematic reviews assessed bracing intervention (rigid and soft orthoses) for idiopathic scoliosis deformity [Table 5]. Two of the reviews were appraised as being of high-quality.^{12,41} The conclusions differed between these two reviews. The authors of the earlier publication (2005) concluded that the effectiveness of bracing is not yet established, but might be promising.¹² In a subsequent systematic review (2015), the conclusions favored bracing, although the evidence was of very low-

*Optum™ Physical Health ("Optum") includes OptumHealth Care Solutions, LLC; ACN Group IPA of New York, Inc.; ACN Group IPA of California, Inc. d/b/a OptumHealth Physical Health of California; Managed Physical Network, Inc.; and OrthoNet Holdings, Inc. which includes OrthoNet New York IPA, Inc., OrthoNet West, Inc., OrthoNet, LLC, OrthoNet of the South, Inc.

quality.⁴¹ This Cochrane review also concluded, that a rigid brace was more successful than an elastic brace at curbing curve progression when measured in Cobb degrees in low degree curves (20° to 30°), with no significant differences in the subjective perception of daily difficulties associated with wearing the brace.⁴¹ The two low-quality literature reviews concurred that the strength of evidence favors bracing over observation.^{42,43}

A small pilot case series (N=5) sought to evaluate the potential benefits of axial spinal unloading – a form of non-operative traction – over a 3-month period.⁴⁴ The authors found reductions in curve magnitude suggesting this therapy may be a potential adjunct in the treatment of adolescent idiopathic scoliosis. The design and shortcomings in the inclusion criteria resulted in very low confidence in the results of this study. The inclusion criteria did not account for a minimum Cobb angle or Risser grade. Two of the five subjects had baseline Cobb angles of <10 degrees. Only one subject had a baseline Cobb angle of >20 degrees.[Table 6]

A modeling study found that scoliotic spines were more sensitive to whole body vibration.⁴⁵ These results suggest that vibration may exacerbate the degree of scoliosis and so such patients should reduce their exposure to vibration.

Research In-Progress:

A search for clinical trials in varying stage of progress (e.g., recruiting, active, complete but not published) identified 106 registered studies (www.clinicaltrials.gov) concerning conservative interventions, predominantly exercise or bracing for idiopathic scoliosis.

Evidence-Informed Practice

Clinical decision support resources have described evidence-informed guidance for the conservative management of individuals diagnosed with idiopathic scoliosis in clinical care settings.^{46,47} Bracing, using rigid orthoses, is viewed as the only active nonsurgical intervention described as proven effective.

UpToDate[®], a point-of-care evidence synthesis resource, has concluded that rigid bracing is a valid option for the treatment of adolescent idiopathic scoliosis in patients meeting established selection criteria.⁴⁶ UpToDate[®] has also determined, “There is a lack of high-quality evidence from randomized trials that chiropractic treatment, electrical stimulation, or biofeedback is effective... Although some randomized trials suggest that physiotherapeutic scoliosis-specific exercises may reduce the Cobb angle and improve health-related quality of life, systematic reviews of randomized and controlled clinical trials have concluded that the benefits are unproven.”

DynaMed, another point-of-care evidence synthesis resource, has arrived at similar conclusions. In patients with adolescent idiopathic scoliosis, bracing is the primary treatment and appears effective. Scoliosis-specific exercises have limited evidence for improving spine curvature or reducing progression of adolescent idiopathic scoliosis.⁴⁷

Table 3: Quality Appraisal of Systematic Reviews for Manual Therapy (including manipulation)

Appraisal Methods		Quality Appraisal				
Item	Description	Everett (2007)	Romano (2008)	Gleberzon (2012)	Posadski (2013)	Theroux (2017)
Was an 'a priori' design provided?	The research question and inclusion criteria should be established before the conduct of the review.	Yes	Yes	Yes	Yes	Yes
Was there duplicate study selection and data extraction?	There should be at least two independent data extractors and a consensus procedure for disagreements should be in place.	Yes	No	Yes	Yes	Yes
Was a comprehensive literature search performed?	At least two electronic sources should be searched. The report must include years and databases used (e.g., Central, EMBASE, and MEDLINE). Key words and/or MESH terms must be stated and where feasible the search strategy should be provided. All searches should be supplemented by consulting current contents, reviews, textbooks, specialized registers, or experts in the particular field of study, and by reviewing the references in the studies found.	Yes	Yes	Yes	Yes	Yes
Was the status of publication (e.g., grey literature) used as an inclusion criterion?	The authors should state that they searched for reports regardless of their publication type. The authors should state whether or not they excluded any reports (from the systematic review), based on their publication status, language etc.	Yes	No	Yes	Yes	Yes
Was a list of studies (included and excluded) provided?	A list of included and excluded studies should be provided.	No	No	No	No	Included-only
Were the characteristics of the included studies provided?	In an aggregated form such as a table, data from the original studies should be provided on the participants, interventions and outcomes. The ranges of characteristics in all the studies analyzed e.g., age, race, sex, relevant socioeconomic data, disease status, duration, severity, or other diseases should be reported.	No	N/A (None of the studies met inclusion criteria)	Yes	Yes	Yes
Was the scientific quality of the included studies assessed and documented?	'A priori' methods of assessment should be provided (e.g., for effectiveness studies if the author(s) chose to include only randomized, double-blind, placebo-controlled studies, or allocation concealment as inclusion criteria); for other types of studies alternative items will be relevant.	No (level of evidence scale only)	N/A	Yes (Sackett score)	Yes (Cochrane Risk of Bias)	Yes (Cochrane Risk of Bias; Effective Practice and Organisation of Care's criteria for nonrandomized Trials)
Was the scientific quality of the included studies used appropriately in formulating conclusions?	The results of the methodological rigor and scientific quality should be considered in the analysis and the conclusions of the review, and explicitly stated in formulating recommendations.	N/A	N/A	Yes	Yes	Yes
Were the methods used to combine the findings of studies appropriate?	For the pooled results, a test should be done to ensure the studies were combinable, to assess their homogeneity (i.e., Chi-squared test for homogeneity, P). If heterogeneity exists a random effects model should be used and/or the clinical appropriateness of combining should be taken into consideration (i.e., is it sensible to combine?).	N/A	N/A	N/A	N/A (single study met inclusion criteria)	N/A
Was the likelihood of publication bias assessed?	An assessment of publication bias should include a combination of graphical aids (e.g., funnel plot, other available tests) and/or statistical tests (e.g., Egger regression test).	No	N/A	No	No	No
Was the conflict of interest stated?	Potential sources of support should be clearly acknowledged in both the systematic review and the included studies.	Yes	N/A	No	Yes	Yes
Score		5	2	7	8	9

*Optum™ Physical Health ("Optum") includes OptumHealth Care Solutions, LLC; ACN Group IPA of New York, Inc.; ACN Group IPA of California, Inc. d/b/a OptumHealth Physical Health of California; Managed Physical Network, Inc.; and OrthoNet Holdings, Inc. which includes OrthoNet New York IPA, Inc., OrthoNet West, Inc., OrthoNet, LLC, OrthoNet of the South, Inc.

Table 4: Quality Appraisal of Systematic Reviews for Exercises Using AMSTAR 2*

Quality Assessment Item	Burger (2019)	Day (2019)	Farooqui (2018)	Laita (2018)	Thompson (2019)	Fan (2020)	Gámiz-Bermúdez (2021)
1. Did the research questions and inclusion criteria for the review include the components of PICO?	Yes	Yes	Yes	Yes	Yes	Yes	Yes
2. Did the report of the review contain an explicit statement that the review methods were established prior to the conduct of the review and did the report justify any significant deviations from the protocol?	No	No	No	No	Partial Yes	Yes	Yes
3. Did the review authors explain their selection of the study designs for inclusion in the review?	Yes	No	No	Yes	Yes	Yes	Yes
4. Did the review authors use a comprehensive literature search strategy?	Partial Yes	Partial Yes	Partial Yes ⁷	Partial Yes	Partial Yes	No ¹³	Partial Yes
5. Did the review authors perform study selection in duplicate?	Yes	No	No	Yes	Yes	Yes	Yes
6. Did the review authors perform data extraction in duplicate?	Yes	No	No	Yes	Yes	Yes	Yes
7. Did the review authors provide a list of excluded studies and justify the exclusions?	No	Yes	No	Yes	Yes	No	No
8. Did the review authors describe the included studies in adequate detail?	Yes	Partial Yes	Partial Yes	Partial Yes	Partial Yes	Partial Yes	Partial Yes
9. Did the review authors use a satisfactory technique for assessing the RoB in individual RCTs that were included in the review?	Yes ¹	Yes	Yes	Yes	Yes	Yes	Yes
10. Did the review authors use a satisfactory technique for assessing the RoB in individual NRSI that were included in the review?	N/A	No ⁵	N/A	N/A	N/A	No ¹⁴	N/A
11. Did the review authors report on the sources of funding for the studies included in the review?	No ²	No	No	No	No	No	No
12. If meta-analysis was performed did the review authors use appropriate methods for statistical combination of results for RCTs?	No ³	No ⁶	No ⁸	N/A	Yes	N/A	No ¹⁶
13. If meta-analysis was performed did the review authors use appropriate methods for statistical combination of results for NRSI?	N/A	No ⁶	N/A	N/A	N/A	N/A	N/A
14. If meta-analysis was performed, did the review authors assess the potential impact of RoB in individual studies on the results of the meta-analysis or other evidence synthesis?	N/A	N/A	No	N/A	N/A ¹¹	N/A	N/A ¹⁷
15. Did the review authors account for RoB in individual studies when interpreting/discussing the results of the review?	Yes	No	No	Yes ⁹	Yes	Yes	Yes
16. Did the review authors provide a satisfactory explanation for, and discussion of, any heterogeneity observed in the results of the review?	No	No	No	No	No	No ¹⁵	Yes
17. If they performed quantitative synthesis did the review authors carry out an adequate investigation of publication bias (small study bias) and discuss its likely impact on the results of the review?	N/A ⁴	No	No	N/A	N/A ¹²	N/A	Yes
18. Did the review authors report any potential sources of conflict of interest, including any funding they received for conducting the review?	Yes	Yes	No	Yes	Yes	Yes	Yes
Rating overall confidence in the results of the review	Low	Low	Critically Low	Low¹⁰	Moderate	Moderate	Low

*Optum™ Physical Health (“Optum”) includes OptumHealth Care Solutions, LLC; ACN Group IPA of New York, Inc.; ACN Group IPA of California, Inc. d/b/a OptumHealth Physical Health of California; Managed Physical Network, Inc.; and OrthoNet Holdings, Inc. which includes OrthoNet New York IPA, Inc., OrthoNet West, Inc., OrthoNet, LLC, OrthoNet of the South, Inc.

* Source: Shea BJ, Reeves BC, Wells G, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ*. 2017 Sep 21;358:j4008.

Legend: PICO = population, intervention, comparator, outcome; MD = mean difference; N/A = not applicable; NRSI = nonrandomized studies of an intervention; RCT = randomized controlled trial; RoB = risk of bias; SMD = standardized mean difference

Rating overall confidence in the results of the review

High	Zero or one non-critical weakness: The systematic review provides an accurate and comprehensive summary of the results of the available studies that address the question of interest.
Moderate	More than one non-critical weakness: The systematic review has more than one weakness, but no critical flaws. It may provide an accurate summary of the results of the available studies that were included in the review.
Low	One critical flaw with or without non-critical weaknesses: The review has a critical flaw and may not provide an accurate and comprehensive summary of the available studies that address the question of interest.
Critically low	More than one critical flaw with or without non-critical weaknesses: The review has more than one critical flaw and should not be relied on to provide an accurate and comprehensive summary of the available studies.

Note: Multiple non-critical weaknesses may diminish confidence in the review and it may be appropriate to move the overall appraisal down from moderate to low confidence

Rationale:

1. Only the RoB summary assessment scores were provided. A table showing the individually assessed items was not included in the study or as supplemental material. The narrative did add information about lack of blinding (participants, clinicians, assessors). [non-critical weakness]
2. Fixed effects meta-analytic model was used when a random effects analysis was the appropriate choice (Figure 2). This likely resulted in overestimating the pooled the effect size (that is, to a more substantial benefit). [critical flaw]
3. Only two studies per meta-analysis
4. Too few studies to assess for publication bias
5. The PEDro checklist for RCTs was used for NRSI
6. Forest plots calculated effect sizes (Hedge's d), when calculations using direct measurement (degrees) would have been more informative e.g., interpreting clinical relevance [critical flaw]. Meta-analysis not performed due to significant heterogeneity.
7. The search strategy included 3 data bases, only 1 of which was a standardized bibliographic platform. [critical flaw]
8. The Q test and I² test were reportedly calculated to identify the level of heterogeneity; however, the results were not described. Further, there was no exploration of the causes of heterogeneity. The authors reported both fixed and random pooled effects (which showed different results). Without knowing the significance of heterogeneity, it was not possible to determine the most appropriate analytic model. The meta-analysis used SMD when MD (degrees) would have been more informative. [critical flaw]
9. The analysis included studies deemed to have a low or moderate risk of bias
10. The analysis reported on statistical significance (p values) but did not provide information about clinically meaningful change (>5°) with intervention [critical flaw]
11. Six of 7 studies had a high RoB
12. Too few studies to conduct an analysis for the potential of publication bias
13. The literature search did not include an explanation for limiting study selection to English language (non-critical flaw)
14. The PEDro checklist was misapplied to NRSI (non-critical flaw)
15. The reviewers did not discuss the heterogeneity of results (non-critical flaw)
16. The effect estimates were statistically reported as SMD, when MD would have been more appropriate. Consequently, the clinical relevance of the results could not be determined (critical flaw)
17. All studies were rated as having a high RoB

Table 5: Quality Appraisal of Systematic Reviews for Bracing Intervention

Appraisal Methods		Quality Appraisal			
Item	Description	Lenssinck (2005)	Maruyama (2011)	Weiss (2012)	Negrini (2015)
Was an 'a priori' design provided?	The research question and inclusion criteria should be established before the conduct of the review.	Yes	Yes	Yes	Yes
Was there duplicate study selection and data extraction?	There should be at least two independent data extractors and a consensus procedure for disagreements should be in place.	Yes	?	?	Yes
Was a comprehensive literature search performed?	At least two electronic sources should be searched. The report must include years and databases used (e.g. Central, EMBASE, and MEDLINE). Key words and/or MESH terms must be stated and where feasible the search strategy should be provided. All searches should be supplemented by consulting current contents, reviews, textbooks, specialized registers, or experts in the particular field of study, and by reviewing the references in the studies found.	Yes	No (only a single data base was queried)	No (only a single data base was queried)	Yes
Was the status of publication (e.g., grey literature) used as an inclusion criterion?	The authors should state that they searched for reports regardless of their publication type. The authors should state whether or not they excluded any reports (from the systematic review), based on their publication status, language etc.	Yes	No	No	Yes
Was a list of studies (included and excluded) provided?	A list of included and excluded studies should be provided.	No	No	No	Yes
Were the characteristics of the included studies provided?	In an aggregated form such as a table, data from the original studies should be provided on the participants, interventions and outcomes. The ranges of characteristics in all the studies analyzed e.g. age, race, sex, relevant socioeconomic data, disease status, duration, severity, or other diseases should be reported.	Yes	Yes	No	Yes
Was the scientific quality of the included studies assessed and documented?	'A priori' methods of assessment should be provided (e.g., for effectiveness studies if the author(s) chose to include only randomized, double-blind, placebo controlled studies, or allocation concealment as inclusion criteria); for other types of studies alternative items will be relevant.	Yes	Yes (Cochrane Risk of Bias)	No	Yes (Cochrane Risk of Bias; Newcastle-Ottawa Scale)
Was the scientific quality of the included studies used appropriately in formulating conclusions?	The results of the methodological rigor and scientific quality should be considered in the analysis and the conclusions of the review, and explicitly stated in formulating recommendations.	Yes	Yes	N/A	Yes
Were the methods used to combine the findings of studies appropriate?	For the pooled results, a test should be done to ensure the studies were combinable, to assess their homogeneity (i.e. Chi-squared test for homogeneity, I ²). If heterogeneity exists a random effects model should be used and/or the clinical appropriateness of combining should be taken into consideration (i.e. is it sensible to combine?).	N/A	N/A	N/A	N/A
Was the likelihood of publication bias assessed?	An assessment of publication bias should include a combination of graphical aids (e.g., funnel plot, other available tests) and/or statistical tests (e.g., Egger regression test).	No	No	No	No
Was the conflict of interest stated?	Potential sources of support should be clearly acknowledged in both the systematic review and the included studies.	No	Yes	Yes	Yes
Score		7	5	2	9

*Optum™ Physical Health (“Optum”) includes OptumHealth Care Solutions, LLC; ACN Group IPA of New York, Inc.; ACN Group IPA of California, Inc. d/b/a OptumHealth Physical Health of California; Managed Physical Network, Inc.; and OrthoNet Holdings, Inc. which includes OrthoNet New York IPA, Inc., OrthoNet West, Inc., OrthoNet, LLC, OrthoNet of the South, Inc.

Table 6. Risk of Bias (limitations in study design or implementation)

Title: Chromy CA, Carey MT, Balgaard KG, Iaizzo PA. The potential use of axial spinal unloading in the treatment of adolescent idiopathic scoliosis: A case series. *Arch Phys Med Rehabil.* 2006;87:1447–1453.

Clear study objective/question	Yes	To evaluate the potential benefits of axial spinal unloading over a 3-month period
Well-defined study protocol	Yes	Protocol included a training session and a pre-formatted “compliance” journal
Explicit inclusion and exclusion criteria for study participants	Yes	Criteria did not account for a minimum Cobb angle or Risser sign. Two of the five subjects had baseline Cobb angles of <10 degrees. Only one subject had a baseline Cobb angle of >20 degrees.*
Specified time interval for patient recruitment	Yes	6-months
Consecutive patient enrollment	Unsure*	Not reported
Clinically relevant outcomes	Yes	
Prospective outcome data collection	Yes	
High follow-up rate	Yes	100%

* = High risk of selection bias

Source: Chan K, Bhandari M. Three-minute critical appraisal of a case series article. *Indian Journal of Orthopaedics* 2011; 45:103–104

References

1. Lenke L (chair). SRS terminology committee and working group on spinal classification revised glossary of terms. *Scoliosis Research Society*. Accessed February 2023. [Revised Glossary of Terms | Scoliosis Research Society \(srs.org\)](#)
2. Konieczny MR, Senyurt H, Krauspe R. Epidemiology of adolescent idiopathic scoliosis. *Journal of Children's Orthopaedics* 2012;7:3–9
3. Rowe DE. The Scoliosis Research Society bracing manual: introduction. *Scoliosis Research Society* 2003. Accessed February 2023. [SRS Bracing Manual | Scoliosis Research Society](#)
4. Thompson GH, Richards BS. Inclusion and assessment criteria for conservative scoliosis treatment. In: Grivas TB, ed. *The Conservative Scoliosis Treatment*. IOS Press, 2008;Section VI:15–163
5. Weiss HR, Goodall D. The treatment of adolescent idiopathic scoliosis (AIS) according to the best evidence: a systematic review. *European Journal of Physical and Rehabilitation Medicine* 2008;44:177–193
6. Canavese F, Kaelin A. Adolescent idiopathic scoliosis: Indications and efficacy of nonoperative treatment. *Indian J Orthop*. 2011; 45: 7–14
7. Weiss H-R, Negrini S, Hawes MC, Rigo M, Kotwicki T, Grivas TB, et al. Physical exercises in the treatment of idiopathic scoliosis at risk of brace treatment -- SOSORT consensus paper 2005. *Scoliosis*. 2006;1:6.
8. Negrini S, Donzelli S, Aulisa AG, et al. 2016 SOSORT guidelines: orthopaedic and rehabilitation treatment of idiopathic scoliosis during growth. *Scoliosis and Spinal Disorders* 2018;13:3
9. Negrini S, Aulisa AG, Aulisa L, et al. 2011 SOSORT guidelines: Orthopaedic and Rehabilitation treatment of idiopathic scoliosis during growth. *Scoliosis* 2012;7:3 doi:10.1186/1748-7161-7-3
10. Monticone M, Ambrosini E, Cazzaniga D, Rocca B, Ferrante S. Active self-correction and task-oriented exercises reduce spinal deformity and improve quality of life in subjects with mild adolescent idiopathic scoliosis. Results of a randomised controlled trial. *Eur Spine J Off Publ Eur Spine Soc Eur Spinal Deform Soc Eur Sect Cerv Spine Res Soc*. 2014;23(6):1204–14.
11. Anwer S, Alghadir A, Shaphe A, Anwar D. Effects of exercise on spinal deformities and quality of life in patients with adolescent idiopathic scoliosis. *BioMed Research International* 2015;2015.
12. Lenssick ML, Frijlink AC, Berger MY, et al. Effect of bracing and other conservative interventions in the treatment of idiopathic scoliosis in adolescents: a systematic review of clinical trials. *Phys Ther* 2005;85:1329–39
13. Stone B, Beekman C, Hall V, Guess V, Brooks HL. The effect of an exercise program on change in curve in adolescents with minimal idiopathic scoliosis. A preliminary study. *Phys Ther*. 1979;59(6):759–63.
14. Wong MS, Cheng JCY, Wong MW, So SF. A work study of the CAD/CAM method and conventional manual method in the fabrication of spinal orthoses for patients with adolescent idiopathic scoliosis. *Prosthetics Orthot Int*. 2005;29(1):93–104.
15. Périé D, Aubin CE, Petit Y, Labelle H, Dansereau J. Personalized biomechanical simulations of orthotic treatment in idiopathic scoliosis. *Clin Biomech Bristol Avon*. 2004;19(2):190–5.
16. Weiss HR, Negrini S, Rigo M, et al. Indications for conservative management of scoliosis (guidelines). *Scoliosis* 2006;1:5 doi:10.1186/1748-7161-1-5
17. Diab AA. The role of forward head correction in management of adolescent idiopathic scoliotic patients: a randomized controlled trial. *Clin Rehabil*. 2012;26(12):1123–32.
18. Toledo PCV, de Mello DB, Araújo ME, Daoud R, Dantas EHM. Global posture reeducation effects in students with scoliosis. *Fisioter E Pesqui*. 2011;18(4):329–34.
19. McMaster MJ. Infantile idiopathic scoliosis: can it be prevented? *J Bone Joint Surg Br*. 1983;65(5):612–7.
20. Higgins JPT, Thomas J, Chandler J, et al (editors). *Cochrane handbook for systematic reviews of interventions version 6.0 (updated July 2019) [Chapter 4: Searching for and selecting studies]*. Cochrane, 2019. Accessed February 2023. www.training.cochrane.org/handbook.
21. Shea BJ, Hamel C, Wells GA, et al. AMSTAR is a reliable and valid measurement tool to assess the methodological quality of systematic reviews. *Journal of Clinical Epidemiology* 2009; 62:1013-1020.
22. Shea BJ, Bouter LM, Peterson J, et al. External Validation of a Measurement Tool to Assess Systematic Reviews (AMSTAR). *PLoS ONE* 2007; 2:e1350.
23. Mikton C, Butchart A. Child maltreatment prevention: a systematic review of reviews. *Bull World Health Organ* 2009; 87:353–361
24. Rookmoneea M, Dennis L, Brealey S, et al. The effectiveness of interventions in the management of patients with primary frozen shoulder. *J Bone Joint Surg [Br]* 2010; 92-B:1267–72
25. Sakzewski L, Ziviani J, Boyd R. Systematic review and meta-analysis of therapeutic management of upper-limb dysfunction in children with congenital hemiplegia. *Pediatrics* 2009; 123:e1111-e1122

*Optum™ Physical Health (“Optum”) includes OptumHealth Care Solutions, LLC; ACN Group IPA of New York, Inc.; ACN Group IPA of California, Inc. d/b/a OptumHealth Physical Health of California; Managed Physical Network, Inc.; and OrthoNet Holdings, Inc. which includes OrthoNet New York IPA, Inc., OrthoNet West, Inc., OrthoNet, LLC, OrthoNet of the South, Inc.

26. Shea BJ, Reeves BC, Wells G, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *British Medical Journal* 2017;358:j4008.
27. Gleberzon BJ, Arts J, Mei A, McManus EL. The use of spinal manipulative therapy for pediatric health conditions: a systematic review of the literature. *J Can Chiropr Assoc.* 2012;56:128–141
28. Posadzki P, Lee MS, Ernst E. Osteopathic manipulative treatment for pediatric conditions: a systematic review. *Pediatrics* 2013;132:140-52. doi: 10.1542/peds.2012-3959.
29. Théroux J, Stomski N, Losco CD, et al. Spinal manipulative therapy for adolescent idiopathic scoliosis: a systematic review. *Journal of Manipulative & Physiological Therapeutics* 2017;40(6):452-8.
30. Czaprowski D. Manual therapy in the treatment of idiopathic scoliosis. Analysis of current knowledge. *Ortopedia, Traumatologia, Rehabilitacja* 2016;18(5):409-424.
31. Lotan S, Kalichman L. Manual therapy treatment for adolescent idiopathic scoliosis. *Journal of Bodywork and Movement Therapies* 2019;23:189-193.
32. Fan Y, Ren Q, To MK, Cheung JP. Effectiveness of scoliosis-specific exercises for alleviating adolescent idiopathic scoliosis: a systematic review. *BMC Musculoskeletal Disorders* 2020 Dec;21(1):1-3.
33. Thompson JY, Williamson EM, Williams MA, Heine PJ, Lamb SE, ACTiVATeS Study Group. Effectiveness of scoliosis-specific exercises for adolescent idiopathic scoliosis compared with other non-surgical interventions: a systematic review and meta-analysis. *Physiotherapy.* 2019 Jun 1;105(2):214-34.
34. Burger M, Coetzee W, du Plessis LZ, et al. The effectiveness of Schroth exercises in adolescents with idiopathic scoliosis: A systematic review and meta-analysis. *The South African Journal of Physiotherapy* 2019;75(1) a904. Accessed February 2023. <https://doi.org/10.4102/sajp.v75i1.904>
35. Day JM, Fletcher J, Coghlan M, Ravine T. Review of scoliosis-specific exercise methods used to correct adolescent idiopathic scoliosis. *Archives of Physiotherapy* 2019;9(1):8.
36. Farooqui SI, Siddiqui PQ, Basit Ansari AF. Effects of spinal mobilization techniques in the management of adolescent idiopathic scoliosis-A meta-analysis. *International Journal of Health Sciences* 2018;12(6):44.
37. Laita LC, Kinesiologista CT, Kinesiologistb TM, del Barrio SJ. Effects of corrective, therapeutic exercise techniques on adolescent idiopathic scoliosis. A systematic review. *Archivos Argentinos de Pediatría* 2018;116(4):e582-9.
38. Gámiz-Bermúdez F, Obrero-Gaitán E, Zagalaz-Anula N, Lomas-Vega R. Corrective exercise-based therapy for adolescent idiopathic scoliosis: Systematic review and meta-analysis. *Clinical Rehabilitation.* 2021 Dec 28:02692155211070452.
39. Li X, Shen J, Liang J, et al. Effect of core-based exercise in people with scoliosis: A systematic review and meta-analysis. *Clinical Rehabilitation.* 2021 May;35(5):669-80.
40. Zhou Z, Liu F, Li R, Chen X. The effects of exercise therapy on adolescent idiopathic scoliosis: an overview of systematic reviews and meta-analyses. *Complementary Therapies in Medicine.* 2021 Feb 23:102697.
41. Negrini S, Minozzi S, Bettany-Saltikov J, et al. Braces for idiopathic scoliosis in adolescents. *Cochrane Database of Systematic Reviews* 2015, Issue 6. Art. No.: CD006850. DOI: 10.1002/14651858.CD006850.pub3.
42. Maruyama T, Grivas TB, Kaspiris A. Effectiveness and outcomes of brace treatment: a systematic review. *Physiotherapy Theory and Practice* 2011;27:26–42
43. Weiss HR, Werkmann M. Soft braces in the treatment of Adolescent Idiopathic Scoliosis (AIS) - Review of the literature and description of a new approach. *Scoliosis* 2012;7:11. doi: 10.1186/1748-7161-7-11.
44. Chromy CA, Carey MT, Balgaard KG, Iazzo PA. The potential use of axial spinal unloading in the treatment of adolescent idiopathic scoliosis: A case series. *Arch Phys Med Rehabil.* 2006;87:1447–1453.
45. Jia S, Li Y, Xie J, et al. Differential response to vibration of three forms of scoliosis during axial cyclic loading: a finite element study. *BMC Musculoskeletal Disorders.* 2019 Dec 1;20(1):370.
46. Scherl SA, Phillips W, Torchia MM. Adolescent idiopathic scoliosis: management and prognosis. UpToDate® Literature review current through Jan. 2023. | This topic last updated: Jan. 4, 2023. Accessed February 2023. https://www.uptodate.com/contents/adolescent-idiopathic-scoliosis-management-and-prognosis?search=scoliosis&source=search_result&selectedTitle=3~150&usage_type=default&display_rank=3
47. DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T116647, Adolescent Idiopathic Scoliosis; [updated Nov. 30, 2018. Accessed February 2023 Available from [DynaMed](#)



Utilization Management Policy

Policy History/Revision Information

Date	Action/Description
1/1997	Original effective date
3/24/1998	Annual review completed
1/28/1999	Annual review completed
2/23/2000	Annual review completed
3/07/2001	Annual review completed
9/04/2001	Annual review completed
2/14/2002	Policy inactivated
4/17/2014	Policy revised: Scope changed to focus on conservative interventions; Methodology conducted in accordance with guidance recorded in policy 429
4/16/2015	Annual review completed
4/21/2016	Annual review completed
4/20/2017	Annual review completed; Legal entity name changed from “OptumHealth Care Solutions, Inc.” to “OptumHealth Care Solutions, LLC.”
4/26/2018	Annual review completed; Literature Review and References revised; Appendix deleted
4/25/2019	Annual review completed. Updated the literature review and references. Plain Language Summary added to the document.
4/23/2020	Annual review completed. Updated the literature review and references. Table 4 was revised using the AMSTAR 2 instrument
4/22/2021	Annual review and approval completed
5/03/2022	Annual review completed. Updated the literature review and evidence-informed practice sections, Table 4, and references
6/29/2022	Updated legal entity name “OptumHealth Care Solutions, LLC.” to *Optum™ Physical Health (“Optum”) includes OptumHealth Care Solutions, LLC; ACN Group IPA of New York, Inc.; ACN Group IPA of California, Inc. d/b/a OptumHealth Physical Health of California; Managed Physical Network, Inc.; and OrthoNet Holdings, Inc. which includes OrthoNet New York IPA, Inc., OrthoNet West, Inc., OrthoNet, LLC, OrthoNet of the South, Inc.
4/27/23	Annual review and approval completed; no significant changes made to the document. Updated contact email from policy.inquiry@optumhealth.com to phpolicy_inquiry@optum.com .

Contact Information

Please forward any commentary or feedback on OptumHealth Care Solutions, LLC (OptumHealth) Utilization Management policies to: phpolicy_inquiry@optum.com with the word “Policy” in the subject line.

The services described in Optum* by OptumHealth Care Solutions, LLC policies are subject to the terms, conditions and limitations of the Member's contract or certificate. OptumHealth reserves the right, in its sole discretion, to modify policies as necessary without prior written notice unless otherwise required by Optum's administrative procedures.

Certain internal policies may not be applicable to self-funded members and certain insured products. Refer to the member's Summary Plan Description (SPD) or Certificate of Coverage (COC) to determine whether coverage is provided or if there are any exclusions or benefit limitations applicable to any of these policies. If there is a difference between any policy and the member's SPD or COC, the member's SPD or COC will govern.

*Optum™ Physical Health (“Optum”) includes OptumHealth Care Solutions, LLC; ACN Group IPA of New York, Inc.; ACN Group IPA of California, Inc. d/b/a OptumHealth Physical Health of California; Managed Physical Network, Inc.; and OrthoNet Holdings, Inc. which includes OrthoNet New York IPA, Inc., OrthoNet West, Inc., OrthoNet, LLC, OrthoNet of the South, Inc.

PLAIN LANGUAGE SUMMARY***Scoliosis: Conservative Interventions*****Utilization Management Policy # 95**

Plain Language Summaries are a service provided by *Optum*^{*} by *OptumHealth Care Solutions, LLC* to help patients better understand the complicated and often mystifying language of modern healthcare.

Plain Language Summaries are presented to supplement the associated clinical policy or guideline. These summaries are not a substitute for advice from your own healthcare provider.

What are conservative interventions for scoliosis and what is known about them so far?

Conservative interventions for scoliosis commonly include bracing, exercises, and manual therapy – a treatment that uses hands-on pressure to gently move your joints and tissues to correct any restrictions in your range of motion.

There is evidence that rigid braces are helpful for preventing or slowing curve progression for adolescents diagnosed with idiopathic scoliosis.

How were conservative interventions for scoliosis evaluated?

A work group of clinicians was assigned to review the available research. The internet was searched for articles about conservative treatments for idiopathic scoliosis. The work group independently examined the selected research studies. A broadly accepted rating scale was used. Possible ratings were high or low quality.

Before it was approved, the policy was presented to a series of committees that included independent health care practitioners.

*Optum[™] Physical Health (“Optum”) includes OptumHealth Care Solutions, LLC; ACN Group IPA of New York, Inc.; ACN Group IPA of California, Inc. d/b/a OptumHealth Physical Health of California; Managed Physical Network, Inc.; and OrthoNet Holdings, Inc. which includes OrthoNet New York IPA, Inc., OrthoNet West, Inc., OrthoNet, LLC, OrthoNet of the South, Inc.

What did the work group find?

The use of a rigid brace appears to be effective at curbing curve progression. Elastic braces are not as effective as rigid braces. There is some evidence showing exercises, including specialized scoliosis exercises, may help with scoliosis curvature. However, additional research is needed before making recommendations. There is too little evidence to make recommendations about the effectiveness of manual therapy for the treatment of curvature associated with scoliosis.

What were the limitations of the information?

The research supporting conservative interventions for idiopathic scoliosis is based upon low quality studies. For the most part, exercise and manual therapy have not been compared to surgery. Additional research will help in more accurately defining the benefit from these services.

What are the conclusions?

Optum considers rigid brace therapy to be proven and medically necessary for the prevention or stabilization of scoliosis curvature.

Soft braces, manual therapy, exercise and other forms of conservative interventions (e.g., traction) are viewed as *unproven and not medically necessary*.